

UC San Diego

UC San Diego Previously Published Works

Title

Crystal structure of dimethyl 9H-carbazole-2,7-di-carb-oxy-late.

Permalink

<https://escholarship.org/uc/item/8c25k37w>

Journal

Acta crystallographica. Section E, Crystallographic communications, 71(Pt 10)

ISSN

2056-9890

Authors

Lehane, Ryan L
Golen, James A
Rheingold, Arnold L
et al.

Publication Date

2015-10-01

DOI

10.1107/s2056989015017557

Peer reviewed



Crystal structure of dimethyl 9H-carbazole-2,7-dicarboxylate

Ryan L. Lehane,^a James A. Golen,^b Arnold L. Rheingold^b and David R. Manke^{a*}

^aDepartment of Chemistry and Biochemistry, University of Massachusetts Dartmouth, 285 Old Westport Road, North Dartmouth, MA 02747, USA, and ^bDepartment of Chemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA. *Correspondence e-mail: dmanke@umassd.edu

Received 9 September 2015; accepted 19 September 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


In the title compound, C₁₆H₁₃NO₄, the carbazole ring system is almost planar with non-H atoms possessing a mean deviation from planarity of 0.037 Å. The two ester groups are orientated *trans* to one another and tilted slightly from the mean plane of the carbazole ring system, making dihedral angles of 8.12 (6) and 8.21 (5)°. In the crystal, molecules are linked by pairs of N—H...O hydrogen bonds forming inversion dimers. The dimers are linked by parallel slipped π – π interactions, forming slabs propagating along the *b*-axis direction [inter-centroid distance = 3.6042 (8) Å, inter-planar distance = 3.3437 (5) Å, slippage = 1.345 Å].

Keywords: crystal structure; carbazoles; hydrogen bonding; π – π interactions.

CCDC reference: 1426074

1. Related literature

For the synthesis of the title compound, see: Olkhovik *et al.* (2008). For the crystal structures of some carbazoles, see: Clarke & Spink (1969); Gajda *et al.* (2014). For the structure of 9H-carbazole-3,6-dicarboxylic acid, see: Weseliński *et al.* (2014). For coordination polymers featuring the dicarboxylate of the parent compound, see: Yi *et al.* (2013, 2014, 2015).



2. Experimental

2.1. Crystal data

| | |
|---|---|
| C ₁₆ H ₁₃ NO ₄ | $V = 2651.2 (3) \text{ \AA}^3$ |
| $M_r = 283.27$ | $Z = 8$ |
| Monoclinic, $C2/c$ | Mo $K\alpha$ radiation |
| $a = 29.684 (2) \text{ \AA}$ | $\mu = 0.10 \text{ mm}^{-1}$ |
| $b = 5.8264 (4) \text{ \AA}$ | $T = 100 \text{ K}$ |
| $c = 15.4210 (11) \text{ \AA}$ | $0.28 \times 0.18 \times 0.10 \text{ mm}$ |
| $\beta = 96.252 (3)^\circ$ | |

2.2. Data collection

| | |
|---|--|
| Bruker CMOS detector | 16278 measured reflections |
| diffractometer | 2716 independent reflections |
| Absorption correction: multi-scan | 2141 reflections with $I > 2\sigma(I)$ |
| (<i>SADABS</i> ; Bruker, 2005) | $R_{\text{int}} = 0.042$ |
| $T_{\text{min}} = 0.972$, $T_{\text{max}} = 0.990$ | |

2.3. Refinement

| | |
|---------------------------------|--|
| $R[F^2 > 2\sigma(F^2)] = 0.037$ | H atoms treated by a mixture of independent and constrained refinement |
| $wR(F^2) = 0.098$ | $\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$ |
| $S = 1.03$ | $\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$ |
| 2716 reflections | |
| 195 parameters | |
| 1 restraint | |

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---------------------|----------|-------------|-------------|---------------|
| $N1-H1N\cdots O3^i$ | 0.87 (1) | 2.04 (1) | 2.8834 (16) | 164 (1) |

Symmetry code: (i) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*, *PLATON* (Spek (2009) and *pubCIF* (Westrip, 2010).

Acknowledgements

RLL thanks the Jean Dreyfus Boissvain Lectureship for Undergraduate Institutions, the UMass Dartmouth Office of Undergraduate Research Award and the UMass Dartmouth Honors Summer Research Grant for funding. DRM gratefully acknowledges support from the National Science Foundation (CHE-1229339 and CHE-1429086).

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU5208).

References

- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clarke, P. T. & Spink, J. M. (1969). *Acta Cryst.* **B25**, 162.
- Gajda, K., Zarychta, B., Kopka, K., Daszkiewicz, Z. & Ejsmont, K. (2014). *Acta Cryst.* **C70**, 987–991.
- Olkhovik, V. K., Vasilevskii, D. A., Pap, A. A., Kalechyts, G. V., Matveienko, Y. V., Baran, A. G., Halinowski, N. A. & Petushok, V. G. (2008). *Arkivoc.* pp. 69–93.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Weseliński, Ł. J., Luebke, R. & Eddaoudi, M. (2014). *Synthesis*, **46**, 596–599.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yi, X.-C., Huang, M.-X., Qi, Y. & Gao, E.-Q. (2014). *Dalton Trans.* **43**, 3691–3697.
- Yi, X.-C., Xi, F.-G., Qi, Y. & Gao, E.-Q. (2015). *RSC Adv.* **5**, 893–900.
- Yi, X.-C., Xi, F.-G., Wang, K., Su, Z. & Gao, E.-Q. (2013). *J. Solid State Chem.* **206**, 293–299.

supporting information

Acta Cryst. (2015). E71, o784–o785 [doi:10.1107/S2056989015017557]

Crystal structure of dimethyl 9*H*-carbazole-2,7-dicarboxylate

Ryan L. Lehane, James A. Golen, Arnold L. Rheingold and David R. Manke

S1. Comment

Biphenyl-4,4'-dicarboxylate and its derivatives are widely used in metal-organic frameworks (MOFs) as linkers. One derivative that is being explored in coordination polymers is 9*H*-carbazole-2,7-dicarboxylate (Yi *et al.*, 2013, 2014, 2015). Herein, we report on the crystal structure of the previously synthesized 9*H*-carbazole-2,7-dicarboxylic acid dimethyl ester (Olkhovik *et al.*, 2008).

The molecular structure of the title compound is shown in Fig. 1. The bond distances and angles are similar to those observed in some carbazole derivatives (Clarke & Spink, 1969; Gajda *et al.*, 2014,) and a structurally characterized carbazole dicarboxylic acid (Weseliński *et al.*, 2014). The carbazole unit is nearly planar with a mean deviation from planarity of 0.037 Å. The carboxylate groups are *trans* to one another and skewed slightly from the mean plane of the carbazole unit, with carbazole-ester dihedral angles of 8.12 (6) and 8.21 (5)°, involving ester groups O1/O2/C3/C13/C14 and O3/O4/C10/C15/C16, respectively.


In the crystal, molecules are linked by a pair of N—H⋯O hydrogen bonds forming inversion dimers (Fig. 2 and Table 1). The dimers are linked by parallel slipped π - π interactions forming slabs propagating along the *b* axis direction [$\text{Cg3}\cdots\text{Cg3}^i = 3.6042$ (8) Å, interplanar distance = 3.3437 (5) Å, slippage 1.345 Å; Cg3 is the centroid of ring C7—C12; symmetry code: (i) $-x+1/2, -y+1/2, -z+1$].

S2. Synthesis and crystallization


The compound was prepared by a literature procedure (Olkhovik *et al.* 2008). Crystals suitable for X-ray diffraction analysis were grown by slow evaporation of a solution in ethanol.

S3. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Hydrogen atom H1N was located in a difference Fourier map and refined with a distance restraint: N—H = 0.87 (2) Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The C-bound H atoms were placed in calculated positions and refined as riding: C—H = 0.95–0.98 Å with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

**Figure 1**

Molecular structure of the title compound, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A view along the *b* axis of the crystal packing of the title compound, with hydrogen bonds shown as dashed lines (see Table 1).

Dimethyl 9H-carbazole-2,7-dicarboxylate*Crystal data*C₁₆H₁₃NO₄ $M_r = 283.27$ Monoclinic, *C2/c*

Hall symbol: -C 2yc

 $a = 29.684 (2) \text{ \AA}$ $b = 5.8264 (4) \text{ \AA}$ $c = 15.4210 (11) \text{ \AA}$ $\beta = 96.252 (3)^\circ$ $V = 2651.2 (3) \text{ \AA}^3$ $Z = 8$ $F(000) = 1184$ $D_x = 1.419 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5867 reflections

 $\theta = 3.1\text{--}26.3^\circ$ $\mu = 0.10 \text{ mm}^{-1}$ $T = 100 \text{ K}$

Block, yellow

 $0.28 \times 0.18 \times 0.10 \text{ mm}$ *Data collection*

Bruker CMOS detector

diffractometer

Radiation source: fine-focus sealed tube

Doubly curved mirrors monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

 $T_{\min} = 0.972$, $T_{\max} = 0.990$

16278 measured reflections

2716 independent reflections

2141 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.042$ $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.7^\circ$ $h = -36 \rightarrow 36$ $k = -7 \rightarrow 7$ $l = -19 \rightarrow 19$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.037$ $wR(F^2) = 0.098$ $S = 1.03$

2716 reflections

195 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0509P)^2 + 1.5495P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.22 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.22 \text{ e \AA}^{-3}$ *Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|----|-------------|--------------|-------------|----------------------------------|
| O1 | 0.46191 (4) | 0.66773 (19) | 0.82835 (7) | 0.0330 (3) |
| O2 | 0.47499 (4) | 0.3231 (2) | 0.88997 (8) | 0.0435 (3) |
| O3 | 0.15036 (3) | 0.56249 (18) | 0.46010 (7) | 0.0281 (3) |

| | | | | |
|------|-------------|--------------|--------------|------------|
| O4 | 0.12301 (3) | 0.23787 (18) | 0.51083 (7) | 0.0287 (3) |
| N1 | 0.31090 (4) | 0.5801 (2) | 0.63776 (8) | 0.0206 (3) |
| H1N | 0.3176 (5) | 0.703 (2) | 0.6095 (10) | 0.025* |
| C1 | 0.34155 (4) | 0.4608 (2) | 0.69505 (8) | 0.0193 (3) |
| C2 | 0.38361 (5) | 0.5315 (2) | 0.73472 (9) | 0.0214 (3) |
| H2A | 0.3958 | 0.6769 | 0.7221 | 0.026* |
| C3 | 0.40714 (5) | 0.3814 (2) | 0.79351 (9) | 0.0224 (3) |
| C4 | 0.38949 (5) | 0.1642 (3) | 0.81051 (10) | 0.0250 (3) |
| H4A | 0.4066 | 0.0629 | 0.8496 | 0.030* |
| C5 | 0.34764 (5) | 0.0959 (2) | 0.77109 (9) | 0.0217 (3) |
| H5A | 0.3358 | −0.0508 | 0.7831 | 0.026* |
| C6 | 0.32293 (5) | 0.2456 (2) | 0.71332 (9) | 0.0185 (3) |
| C7 | 0.27864 (5) | 0.2357 (2) | 0.66416 (8) | 0.0179 (3) |
| C8 | 0.24369 (5) | 0.0734 (2) | 0.65636 (9) | 0.0196 (3) |
| H8A | 0.2472 | −0.0684 | 0.6868 | 0.024* |
| C9 | 0.20400 (5) | 0.1208 (2) | 0.60415 (9) | 0.0195 (3) |
| H9A | 0.1802 | 0.0107 | 0.5982 | 0.023* |
| C10 | 0.19859 (4) | 0.3317 (2) | 0.55960 (9) | 0.0190 (3) |
| C11 | 0.23263 (4) | 0.4955 (2) | 0.56654 (9) | 0.0189 (3) |
| H11A | 0.2287 | 0.6376 | 0.5365 | 0.023* |
| C12 | 0.27267 (4) | 0.4461 (2) | 0.61861 (9) | 0.0183 (3) |
| C13 | 0.45135 (5) | 0.4484 (3) | 0.84215 (10) | 0.0271 (3) |
| C14 | 0.50435 (5) | 0.7491 (3) | 0.87314 (12) | 0.0371 (4) |
| H14A | 0.5076 | 0.9135 | 0.8620 | 0.056* |
| H14B | 0.5296 | 0.6655 | 0.8518 | 0.056* |
| H14C | 0.5046 | 0.7233 | 0.9360 | 0.056* |
| C15 | 0.15579 (5) | 0.3916 (2) | 0.50484 (9) | 0.0205 (3) |
| C16 | 0.08012 (5) | 0.2876 (3) | 0.46017 (11) | 0.0348 (4) |
| H16A | 0.0590 | 0.1609 | 0.4661 | 0.052* |
| H16B | 0.0849 | 0.3058 | 0.3986 | 0.052* |
| H16C | 0.0675 | 0.4298 | 0.4815 | 0.052* |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|------------|------------|------------|-------------|-------------|-------------|
| O1 | 0.0263 (6) | 0.0291 (6) | 0.0403 (7) | −0.0047 (5) | −0.0108 (5) | 0.0021 (5) |
| O2 | 0.0376 (7) | 0.0349 (7) | 0.0522 (8) | 0.0020 (5) | −0.0209 (6) | 0.0067 (6) |
| O3 | 0.0248 (6) | 0.0295 (6) | 0.0291 (6) | −0.0003 (4) | −0.0013 (4) | 0.0094 (5) |
| O4 | 0.0209 (5) | 0.0290 (6) | 0.0346 (6) | −0.0051 (4) | −0.0045 (5) | 0.0051 (5) |
| N1 | 0.0205 (6) | 0.0176 (6) | 0.0229 (6) | −0.0017 (5) | −0.0012 (5) | 0.0058 (5) |
| C1 | 0.0210 (7) | 0.0189 (7) | 0.0182 (7) | 0.0032 (5) | 0.0024 (5) | 0.0010 (5) |
| C2 | 0.0213 (7) | 0.0200 (7) | 0.0227 (7) | 0.0007 (6) | 0.0014 (6) | 0.0001 (6) |
| C3 | 0.0219 (7) | 0.0236 (7) | 0.0212 (7) | 0.0026 (6) | 0.0003 (6) | −0.0012 (6) |
| C4 | 0.0276 (8) | 0.0240 (7) | 0.0225 (7) | 0.0054 (6) | −0.0008 (6) | 0.0038 (6) |
| C5 | 0.0267 (8) | 0.0182 (7) | 0.0201 (7) | 0.0012 (6) | 0.0022 (6) | 0.0022 (6) |
| C6 | 0.0218 (7) | 0.0182 (7) | 0.0157 (7) | 0.0007 (5) | 0.0035 (5) | −0.0022 (5) |
| C7 | 0.0219 (7) | 0.0187 (7) | 0.0135 (6) | 0.0025 (5) | 0.0040 (5) | −0.0010 (5) |
| C8 | 0.0253 (7) | 0.0163 (6) | 0.0180 (7) | 0.0006 (5) | 0.0053 (6) | 0.0011 (5) |

| | | | | | | |
|-----|------------|-------------|-------------|-------------|-------------|-------------|
| C9 | 0.0219 (7) | 0.0175 (7) | 0.0198 (7) | −0.0027 (5) | 0.0046 (6) | −0.0025 (5) |
| C10 | 0.0206 (7) | 0.0200 (7) | 0.0167 (7) | 0.0002 (6) | 0.0036 (5) | −0.0017 (6) |
| C11 | 0.0222 (7) | 0.0165 (6) | 0.0182 (7) | 0.0013 (5) | 0.0034 (5) | 0.0020 (5) |
| C12 | 0.0204 (7) | 0.0173 (6) | 0.0173 (6) | −0.0009 (5) | 0.0032 (5) | −0.0010 (5) |
| C13 | 0.0265 (8) | 0.0267 (8) | 0.0270 (8) | 0.0041 (6) | −0.0025 (6) | −0.0021 (6) |
| C14 | 0.0266 (9) | 0.0372 (9) | 0.0446 (10) | −0.0064 (7) | −0.0093 (7) | −0.0028 (8) |
| C15 | 0.0212 (7) | 0.0213 (7) | 0.0192 (7) | −0.0016 (6) | 0.0031 (6) | −0.0015 (6) |
| C16 | 0.0218 (8) | 0.0428 (10) | 0.0374 (9) | −0.0036 (7) | −0.0068 (7) | 0.0025 (8) |

Geometric parameters (Å, °)

| | | | |
|------------|-------------|---------------|-------------|
| O1—C13 | 1.3382 (19) | C5—H5A | 0.9500 |
| O1—C14 | 1.4490 (18) | C6—C7 | 1.445 (2) |
| O2—C13 | 1.2070 (18) | C7—C8 | 1.3992 (19) |
| O3—C15 | 1.2119 (17) | C7—C12 | 1.4145 (19) |
| O4—C15 | 1.3333 (17) | C8—C9 | 1.381 (2) |
| O4—C16 | 1.4487 (18) | C8—H8A | 0.9500 |
| N1—C12 | 1.3824 (17) | C9—C10 | 1.409 (2) |
| N1—C1 | 1.3842 (17) | C9—H9A | 0.9500 |
| N1—H1N | 0.870 (13) | C10—C11 | 1.3852 (19) |
| C1—C2 | 1.3915 (19) | C10—C15 | 1.4884 (19) |
| C1—C6 | 1.4112 (19) | C11—C12 | 1.3902 (19) |
| C2—C3 | 1.391 (2) | C11—H11A | 0.9500 |
| C2—H2A | 0.9500 | C14—H14A | 0.9800 |
| C3—C4 | 1.406 (2) | C14—H14B | 0.9800 |
| C3—C13 | 1.491 (2) | C14—H14C | 0.9800 |
| C4—C5 | 1.381 (2) | C16—H16A | 0.9800 |
| C4—H4A | 0.9500 | C16—H16B | 0.9800 |
| C5—C6 | 1.3963 (19) | C16—H16C | 0.9800 |
| | | | |
| C13—O1—C14 | 116.30 (12) | C8—C9—H9A | 119.9 |
| C15—O4—C16 | 115.68 (12) | C10—C9—H9A | 119.9 |
| C12—N1—C1 | 108.67 (11) | C11—C10—C9 | 121.32 (13) |
| C12—N1—H1N | 125.6 (10) | C11—C10—C15 | 116.91 (12) |
| C1—N1—H1N | 124.2 (10) | C9—C10—C15 | 121.75 (12) |
| N1—C1—C2 | 128.90 (13) | C10—C11—C12 | 118.22 (12) |
| N1—C1—C6 | 109.20 (12) | C10—C11—H11A | 120.9 |
| C2—C1—C6 | 121.84 (12) | C12—C11—H11A | 120.9 |
| C1—C2—C3 | 117.59 (13) | N1—C12—C11 | 129.57 (12) |
| C1—C2—H2A | 121.2 | N1—C12—C7 | 109.14 (12) |
| C3—C2—H2A | 121.2 | C11—C12—C7 | 121.27 (12) |
| C2—C3—C4 | 121.10 (13) | O2—C13—O1 | 122.98 (14) |
| C2—C3—C13 | 121.01 (13) | O2—C13—C3 | 124.78 (14) |
| C4—C3—C13 | 117.87 (13) | O1—C13—C3 | 112.24 (12) |
| C5—C4—C3 | 120.86 (13) | O1—C14—H14A | 109.5 |
| C5—C4—H4A | 119.6 | O1—C14—H14B | 109.5 |
| C3—C4—H4A | 119.6 | H14A—C14—H14B | 109.5 |
| C4—C5—C6 | 119.08 (13) | O1—C14—H14C | 109.5 |

| | | | |
|---------------|--------------|-----------------|--------------|
| C4—C5—H5A | 120.5 | H14A—C14—H14C | 109.5 |
| C6—C5—H5A | 120.5 | H14B—C14—H14C | 109.5 |
| C5—C6—C1 | 119.49 (13) | O3—C15—O4 | 122.58 (13) |
| C5—C6—C7 | 133.96 (13) | O3—C15—C10 | 124.62 (13) |
| C1—C6—C7 | 106.53 (12) | O4—C15—C10 | 112.79 (12) |
| C8—C7—C12 | 119.44 (13) | O4—C16—H16A | 109.5 |
| C8—C7—C6 | 134.08 (13) | O4—C16—H16B | 109.5 |
| C12—C7—C6 | 106.46 (12) | H16A—C16—H16B | 109.5 |
| C9—C8—C7 | 119.52 (13) | O4—C16—H16C | 109.5 |
| C9—C8—H8A | 120.2 | H16A—C16—H16C | 109.5 |
| C7—C8—H8A | 120.2 | H16B—C16—H16C | 109.5 |
| C8—C9—C10 | 120.24 (13) | | |
| C12—N1—C1—C2 | −177.05 (13) | C8—C9—C10—C15 | −178.18 (12) |
| C12—N1—C1—C6 | 0.08 (15) | C9—C10—C11—C12 | 0.28 (19) |
| N1—C1—C2—C3 | 177.00 (13) | C15—C10—C11—C12 | 178.78 (12) |
| C6—C1—C2—C3 | 0.2 (2) | C1—N1—C12—C11 | 177.91 (13) |
| C1—C2—C3—C4 | 1.5 (2) | C1—N1—C12—C7 | −0.13 (15) |
| C1—C2—C3—C13 | −176.94 (13) | C10—C11—C12—N1 | −178.41 (13) |
| C2—C3—C4—C5 | −1.8 (2) | C10—C11—C12—C7 | −0.58 (19) |
| C13—C3—C4—C5 | 176.68 (13) | C8—C7—C12—N1 | 178.58 (12) |
| C3—C4—C5—C6 | 0.4 (2) | C6—C7—C12—N1 | 0.13 (15) |
| C4—C5—C6—C1 | 1.3 (2) | C8—C7—C12—C11 | 0.35 (19) |
| C4—C5—C6—C7 | −177.33 (14) | C6—C7—C12—C11 | −178.10 (12) |
| N1—C1—C6—C5 | −178.98 (12) | C14—O1—C13—O2 | 1.1 (2) |
| C2—C1—C6—C5 | −1.6 (2) | C14—O1—C13—C3 | 179.87 (13) |
| N1—C1—C6—C7 | −0.01 (14) | C2—C3—C13—O2 | −175.23 (15) |
| C2—C1—C6—C7 | 177.37 (12) | C4—C3—C13—O2 | 6.3 (2) |
| C5—C6—C7—C8 | 0.6 (3) | C2—C3—C13—O1 | 6.0 (2) |
| C1—C6—C7—C8 | −178.20 (14) | C4—C3—C13—O1 | −172.48 (13) |
| C5—C6—C7—C12 | 178.69 (14) | C16—O4—C15—O3 | 0.0 (2) |
| C1—C6—C7—C12 | −0.07 (14) | C16—O4—C15—C10 | 179.05 (12) |
| C12—C7—C8—C9 | 0.20 (19) | C11—C10—C15—O3 | 7.1 (2) |
| C6—C7—C8—C9 | 178.13 (14) | C9—C10—C15—O3 | −174.35 (13) |
| C7—C8—C9—C10 | −0.49 (19) | C11—C10—C15—O4 | −171.85 (12) |
| C8—C9—C10—C11 | 0.3 (2) | C9—C10—C15—O4 | 6.65 (18) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|---------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N1—H1N \cdots O3 ⁱ | 0.87 (1) | 2.04 (1) | 2.8834 (16) | 164 (1) |

Symmetry code: (i) $-x+1/2, -y+3/2, -z+1$.